

Spectral Sequences in String Topology

Lennart Meier

April 6, 2010

Abstract

In this paper, we investigate the behaviour of the Serre spectral sequence with respect to the algebraic structures of string topology in generalized homology theories, specifically with the Chas-Sullivan product and the corresponding coproduct and the module structures. We prove compatibility for two kinds of fibre bundles: the fibre bundle $\Omega^n M \rightarrow L^n M \rightarrow M$ for an h_* -oriented manifold M and the looped fibre bundle $L^n F \rightarrow L^n E \rightarrow L^n B$ of a fibre bundle $F \rightarrow E \rightarrow B$ of h_* -oriented manifolds. Our method lies in the construction of Gysin morphisms of spectral sequences. We apply these results to study the ordinary homology of the free loop spaces of sphere bundles and generalized homologies of the free loop spaces of spheres and projective spaces.

1 Introduction

Let h_* be a homology theory and M be a d -dimensional h_* -oriented smooth manifold. In [C-S], Chas and Sullivan defined a product on the singular homology of the free loop space and Cohen and Jones generalized it in [C-J] to the case of an arbitrary homology theory h_* . This product is now called the *Chas-Sullivan product* and is of the form

$$h_p(LM) \otimes h_q(LM) \rightarrow h_{p+q-d}(LM).$$

Later several people generalized this product, giving a whole bunch of algebraic structures on different mapping spaces between manifolds, e.g. coproduct and module structures.

To enhance the calculational perspectives, it is, of course, useful to understand the behaviour of these algebraic structures in the Serre spectral sequence associated to certain fibre bundles. The first theorem of this kind was proven in [CJY]. We will generalize their theorem to arbitrary homology theories. Denoting the Serre spectral sequence of a fibre bundle ξ by $\mathcal{E}(\xi)$ and by $[(a, b)]$ a bidegree shift, we have

Theorem 4.7. *Let M be a d -dimensional h_* -oriented manifold. Then*

$$\mathcal{E}(\Omega^n M \rightarrow L^n M \rightarrow M)[(-d, 0)]$$

can be equipped with the structure of a multiplicative spectral sequence which converges to the Chas-Sullivan product on $h_(L^n M)$. Furthermore, the induced product on the E^2 -term $H_{*-d}(M; \mathfrak{h}_q(\Omega^n M))$ is equal to the intersection product with coefficients in the local system of rings $\mathfrak{h}_*(\Omega^n M)$ whose multiplication is given by the Pontryagin product.*

Furthermore, we have

Theorem 4.8. *Let $M \rightarrow N \rightarrow O$ be a fibre bundle of h_* -oriented manifolds of dimensions m , n and o respectively, with projection map π . Then*

$$\mathcal{E}(L^n M \rightarrow L^n N \rightarrow L^n O)[(-o, -m)]$$

can be equipped with the structure of a multiplicative spectral sequence which converges to the Chas-Sullivan product on $h_(L^n N)$. Furthermore, the induced product on the E^2 -term $H_{p+m}(L^n O; \mathfrak{h}_{q+m}(L^n M))$ is equal to the Chas-Sullivan product with coefficients in the local system of rings $\mathfrak{h}_{*+m}(L^n O)$.*

There are similar theorems about coproduct and module structures.

We get these theorems as corollaries from the existence of two kinds of Gysin morphisms (i.e. "wrong-way maps") of spectral sequences, which may be of independent interest and lie at the technical heart of this paper.

Theorem 4.1 (Intersection on the Base). *Let ξ be a fibre bundle with a finite-dimensional manifold B as base and fibre F and let $A \subset B$ be a closed submanifold of codimension d with h_* -oriented normal bundle. Then there is a morphism $s_B(A)$ of convergent spectral sequences of bidegree $(-d, 0)$ between the Serre spectral sequences $\mathcal{E}(\xi)$ and $\mathcal{E}(\xi|_A)$ which induces the usual Gysin morphism $H_p(B; \mathfrak{h}_q(F)) \rightarrow H_{p-d}(A; \mathfrak{h}_q(F))$ on E^2 .*

Theorem 4.5 (Intersection in the Fibre). *Let $\xi = (F \rightarrow E \rightarrow B)$ be a smooth fibre bundle with base a finite-dimensional manifold and fibre a Hilbert manifold and let $\xi_0 = (F_0 \rightarrow E_0 \rightarrow B)$ be a subbundle of constant codimension d with h_* -oriented normal bundle. Then there is a morphism $s_F(E_0)$ of convergent spectral sequences of bidegree $(0, -d)$ between the Serre spectral sequences $\mathcal{E}(\xi)$ and $\mathcal{E}(\xi_0)$ which induces the usual Gysin morphism*

$$H_p(B; \mathfrak{h}_q(F)) \rightarrow H_p(B; \mathfrak{h}_{q-d}(F_0))$$

on E^2 and converges to the Gysin morphism in the homology of the total spaces.

The proofs of these results use Jakob's bordism-like description of an arbitrary homology theory, which was introduced into string topology by Chataur in [Cha]. Both results can be generalized to suitable Hilbert manifolds as base.

As in every mathematical discipline, it is crucial to compute and understand examples to fill the abstract definitions with life. First, we want to compute the Chas-Sullivan product for a certain class of sphere bundles over spheres by rational homotopy theory. Then, we give explicit manifold generators for the singular homologies of the free loop spaces of spheres and projective spaces and use this informations to do computations in complex and oriented bordism and in Landweber exact theories. We get complete answers in the case of odd-dimensional spheres.

Most of the theorems about the behaviour of the Serre spectral sequence with respect to the Chas-Sullivan product and other algebraic structures were already shown by other people in the case of singular homology: We already mentioned [CJY]. Le Borgne ([LBo]) has constructed the Gysin morphisms of spectral sequences and applied them in the cited paper and in [LBo2] to something analogous to 4.8 and 4.10. Kallel and Salvatore ([K-S]) have proven compatibility of the Serre spectral sequences associated to $\Omega^n M \rightarrow L^n M \rightarrow M$ with module structures. To the knowledge of the author, all these spectral sequences are

new for other homology theories. It should be noted that the techniques of the mentioned authors do not generalize since they use chain methods.

Le Borgne has also computed homologies of free loop spaces of sphere bundles in some other cases by a different method.

The paper is structured as follows:

In section 2, we discuss some preliminaries. First, we recall the definition of a Hilbert manifold and prove a certain relative transversality result, which might be well-known, but where I was unable to find a precise reference. These notions are important for our project since (Sobolev) mapping spaces between manifolds provide examples of Hilbert manifolds. Then we recall the definition of Jakob's geometric homology and give a short discussion of Gysin morphisms both in the finite and in the infinite-dimensional case.

In section 3, we recall the definition of the Chas-Sullivan product and also of module and coproduct structures, which will be the basic objects of the paper.

In section 4, we construct first the intersection on the base (in the sense of 4.1) in the finite-dimensional case and use then naturality and approximation by finite-dimensional manifolds to generalize it to the infinite-dimensional case. After constructing also the intersection in the fibre, we prove several statements about the behaviour of the Serre spectral sequence with respect to product, coproduct and module structures.

In section 5, we begin by considering free loop spaces of sphere bundles. By rational homotopy theory, we can prove that the Serre spectral sequence collapses at E^2 in many cases. Then we construct very concrete manifold generators of the free loop spaces of spheres and (complex and quaternionic) projective spaces and use these to prove that the Atiyah-Hirzebruch spectral sequence collapses at E^2 for these spaces in various homology theories.

Acknowledgements

Most of the results in this paper are part of my diploma thesis [Mei1], written under the supervision of Matthias Kreck. To him belongs my gratitude for encouraging support and helpful advice.

2 Preliminaries

2.1 Hilbert Manifolds

A *Hilbert manifold* is a metrizable space which is locally homeomorphic to a separable Hilbert space $\mathbb{E}$. One can define smooth Hilbert manifolds and their tangents spaces in analogy to the finite-dimensional case. We will assume all Hilbert manifolds to be smooth in this paper. To define later Gysin morphisms, we begin with certain transversality results. We do not claim originality here as similar, but deeper results were already proven by Quinn in [Qui].

Lemma 2.1. *Let $E \rightarrow M$ be a Euclidean smooth Hilbert space bundle over a compact manifold M , possibly with boundary. Furthermore, let $L_0, L_1, \dots \subset E$ be a countable collection of sub Hilbert manifolds of finite codimension and $\varepsilon: M \rightarrow \mathbb{R}$ a positive function on M . Then there is a smooth section $s: M \rightarrow E$ with $|s(p)| < \varepsilon(p)$ for all $p \in M$ such that s is transverse to all L_i . If $A \subset M$ is a closed submanifold and the zero section is already transverse to the L_i on A , we can choose $s|_A = 0$.*

Proof. The proof is completely analogous to the finite-dimensional case if one uses the corresponding results in differential topology for Hilbert manifolds and Kuiper's theorem that every Hilbert space bundle is trivial. See e.g. [Mei1, section 2.4.3]. $\square$

Theorem 2.2 (Relative Transversality Theorem). *Let $\pi: E \rightarrow B$ be a fibre bundle where the fibre and the base are Hilbert manifolds. Furthermore, let $E_0^0, E_0^1, \dots \subset E$ be a countable collection of subbundles which are in every fibre sub Hilbert manifolds of finite codimension. Let $f: M \rightarrow E$ be a smooth map from a compact manifold M . Then there is a homotopy $H: M \times I \rightarrow E$ between f and a map $g: M \rightarrow E$ which is transverse to all E_0^i such that $\pi \circ H = \pi \circ f \circ \text{pr}_1$. If $A \subset M$ is a closed submanifold with $f|_A$ transverse to all E_0^i , we can choose $H|_{A \times I} = f \circ \text{pr}_1$.*

Remark 2.3. The theorem generalizes immediately to the case where the base is not a Hilbert manifold, but a differentiable space in the sense of Sikorski, see the discussion before 2.31 in [Mei1]. Note also that for $B = pt$, we get simply the usual (absolute) transversality theorem.

Proof. Consider the closed embedding $\text{id} \times f: M \rightarrow M \times_B E$, where the pullback is over $\pi \circ f: M \rightarrow B$. We can construct a tubular neighbourhood T of M in $M \times_B E$ and identify it with a neighbourhood in the normal bundle. Now choose a section $s: M \rightarrow T$ transverse to all $M \times_B E_0^i$. We can identify the tangent bundle of $M \times_B E$ with $\text{pr}_1^* TM \oplus \text{pr}_2^* T_v E$ where $T_v E$ denotes the vertical part of TE . Since $\pi|_{E_0^i}: E_0^i \rightarrow B$ is a submersion for all i , we have that $s: M \rightarrow M \times_B E \rightarrow M \times E$ is transverse to all $M \times E_0^i$. Hence, we get that $g := \text{pr}_2 \circ s$ is transverse to all E_0^i . $\square$

An important example for a Hilbert manifold is the space $H^n(M, N)$ of Sobolev maps between a compact manifold M of dimension n and an arbitrary manifold N , which is homotopy equivalent to $\text{Map}(M, N)$ with the usual compact-open topology. These results are surely well known for a long time, but I was unable to find a complete proof in the literature. A proof can be found in the companion paper [Mei2], where also a precise definition of $H^n(M, N)$ and further references are given. In the following, we will write $\text{Map}(M, N)$ for $H^n(M, N)$, which will be no source of confusion since at the end we are only interested in the homotopy type.

A useful fact about these mapping spaces is the following approximation theorem:

Theorem 2.4. *Let M, N be manifolds and assume M to be compact. Then there exists a sequence of submanifolds $P_1 \subset P_2 \subset \dots \subset \text{Map}(M, N)$ such that one can deform every map $X \rightarrow \text{Map}(M, N)$ from a compact X to a map into one of the P_i .*

This is a generalization of the corresponding well-known theorem for the loop space. A proof along the lines of [Mil, §16], can be found in [Mei1, section 2.6.2].

2.2 Geometric Homology

In this section, we want to give a bordism description for homology due to Martin Jakob ([Jak]), which works for every (generalized) homology theory. It can be thought as a geometric way to build out of a cohomology theory the corresponding homology theory.

Definition 2.5 (Geometric cycles). Let h^* be a cohomology theory and (X, A) a pair of topological spaces. A *geometric cycle* is a triple (P, a, f) where $f: P \rightarrow X$ is a continuous map

from a compact connected h^* -oriented manifold P with boundary to X such that $f(\partial P) \subset A$ and $a \in h^*(P)$.

If P is of dimension p and $a \in h^m(P)$, then (P, a, f) is a geometric cycle of degree $p - m$.

Take the free abelian group generated by all the geometric cycles and impose the following relation:

$$(P, \lambda a + \mu b, f) = \lambda(P, a, f) + \mu(P, b, f).$$

Thus, we get a graded abelian group. In order to recover the usual homology we must impose two additional relations on geometric cycles:

1. (Bordism relation) We call two triples (P, a, f) and (P', a', f') bordant, if there is a geometric cycle (W, b, g) , such that $P \amalg (-P') \subset \partial W$ is a regularly embedded submanifold of codimension 0 which inherits the h^* -orientation of W . We require further that $b|_P = a, b|_{P'} = a', g|_P = f, g|_{P'} = f'$ and $g(\partial W - P \amalg P') \subset A$. Two bordant cycles are defined to be equivalent.
2. (Vector bundle modification) Let (P, a, f) be a geometric cycle and consider a smooth h^* -oriented d -dimensional vector bundle $\pi: E \rightarrow P$, take the unit sphere bundle $S(E \oplus 1)$ of the Whitney sum of E with a copy of the trivial line bundle over P . The bundle $S(E \oplus 1)$ admits a section s . By $s_!: h^*(P) \rightarrow h^{*+d}(S(E \oplus 1))$ we denote the Gysin morphism in cohomology associated to this section¹. We impose that

$$(P, a, f) \sim (S(E \oplus 1), s_!(a), fp).$$

We lay upon the group of cycles the equivalence relations generated by the relations 1 and 2. An equivalence class of geometric cycle is called a *geometric class* and will be denoted by $[P, a, f]$. We define $gh_q(X, A)$ to be the abelian group of geometric classes of degree q .

Theorem 2.6 ([Jak], Corollary 4.3). *There is a natural isomorphism*

$$gh_q(X, A) \rightarrow h_q(X, A)$$

defined by

$$[P, a, f] \mapsto f_*(a \cap [P]),$$

where $[P]$ is the fundamental class of $(P, \partial P)$ and h_* is the homology theory corresponding to the spectrum representing h^* .

We will identify gh_* and h_* via this isomorphism in the rest of this paper. For later applications, we give an explicit description of the excision isomorphism: Let $[P, a, f] \in h_*(X, A)$ be a geometric class and $B \subset A$ such that $\overline{B} \subset \overset{\circ}{A}$. The preimages $f^{-1}(\overline{B})$ and $f^{-1}(X - \overset{\circ}{A})$ are closed and we can choose a smooth Urysohn function $g: P \rightarrow \mathbb{R}$ separating them. Choose a regular value x between 0 and 1. Then $Q := g^{-1}([0, x])$ is a manifold with boundary in $A - B$. The restriction $[Q, a|_Q, f|_Q]$ is the image of the excision isomorphism in $h_*(X - B, A - B)$. Indeed, $(P \times [0, 1], \text{pr}_1^*(a), f \circ \text{pr}_1)$ is a bordism between $[P, a, f]$ and $i_*[Q, a|_Q, f|_Q]$ since $Q \amalg P$ is a regular submanifold of codimension 0 in $P \amalg P$.

¹This can, for example, be defined via Poincare duality. For other possibilites in the analogous case of homology, see the next section.

2.3 Gysin morphisms

For the definition of the Chas-Sullivan product, the construction of Gysin morphisms (also called *umkehr maps* in the literature) will be especially important. For an inclusion $\iota: A \hookrightarrow B$ of a sub Hilbert manifold of finite codimension d with h_* -oriented normal bundle, a Gysin morphism is a "wrong-way" map $\iota^!: h_*(B) \rightarrow h_{*-d}(A)$. We will give two constructions in the general case and a third one in the finite-dimensional case.

The first uses the theory of geometric homology²: Let $[P, a, f]$ be a geometric cycle in $h_p(B)$. By the transversality theorem, we can assume that f is transversal to A . Now, we define $\iota_G^!([P, a, f]) := [\tilde{P}, a|_{\tilde{P}}, f|_{\tilde{P}}] \in h_{p-d}(A)$, where $\tilde{P} := f^{-1}(A)$. It is easy to see that this class is well-defined. This is the description in the infinite-dimensional case we will primarily use in this paper.

Another possible construction uses the Thom isomorphism theorem: Let $\nu: N \rightarrow A$ be the normal bundle of A and $u \in h^d(N, N - A)$ the Thom class. Then the composition

$$h_*(B) \rightarrow h_*(B, B - A) \cong h_*(N, N - A) \xrightarrow{\cap u} h_{*-d}(N) \cong h_{*-d}(A)$$

is an alternative way to define Gysin maps. It coincides with the construction above in the finite-dimensional case and also for mapping spaces between manifolds by the approximation theorem 2.4. Therefore, our definition of the Chas-Sullivan product will agree with that of [C-J] and [C-V]. Note that one can substitute for h_* also ordinary homology with *local* coefficients.

In the finite-dimensional case, there is also a cellular construction of Gysin maps. To make this precise, we need the following (simple) lemma which is proven in [Mei1, section 2.5]:

Lemma 2.7. *Let $A \subset B$ be a closed submanifold of a finite-dimensional manifold. Then one can triangulate B transversal to A in the sense that every stratum is transverse to A . Furthermore, one can triangulate every $A \cap \Delta_i$ for the simplices Δ_i of $\mathcal{T}$ in a way such that the triangulations coincide in the intersections $A \cap \Delta_i \cap \Delta_j$. Thus one obtains a induced triangulation of A .*

Let $C_*(A)$ and $C_*(B)$ denote the cellular chain complexes. By sending a simplex Δ of the triangulation of B to the sum of all the simplices of $\Delta \cap A$, we get a chain map $s: C_*(B) \rightarrow C_{*-d}(A)$ which induces a Gysin map on homology.

This cellular description of the Gysin morphism is also suitable to describe Gysin morphisms for homology with local coefficients: Let $\mathcal{G}$ be a local system and $x = x_\Delta$ be the midpoint of a simplex Δ . In the cellular complex with respect to $\mathcal{G}$ the coefficient of Δ lies in $\mathcal{G}_x$. Choose arbitrary paths from x to the midpoints of the simplices of $\Delta \cap A$ in Δ and map via them the coefficient of Δ to coefficients for these simplices (note that all possible choices of these paths are homotopic). This describes a Gysin map $g^!: H_*(B; \mathcal{G}) \rightarrow H_{*-d}(A; \mathcal{G}|_A)$.

We want to show the equivalence of the cellular construction with the construction via the Thom isomorphism in the case of singular homology. We can form the sub chain complex $T_*(N, N - A)$ of all singular chains $S_*(N, N - A)$ which are transverse to A (i.e. transverse in each stratum). Given a cycle in $S_*(N, N - A)$, one can form an associated simplicial complex K with a map $f: K \rightarrow B$ by glueing the simplices. By the transversality theorem,

²We follow here [Cha].

one can homotope f to a map g transverse to A without moving ∂K into A . Then (K, g) defines a cycle in $T_*(N, N - A)$ which is homologous to (K, f) . A similar argument can be applied for boundaries. Therefore, $T_*(N, N - A)$ is quasi-isomorphic to $S_*(N, N - A)$. By the universal coefficient theorem, the dual chain complexes are also quasi-isomorphic. Therefore, it is enough to define the Thom class $u \in H^d(N, N - A)$ on d -simplices transverse to A .

For a transverse d -simplex (Δ, f) , we define $u(\Delta)$ as the oriented intersection number $\Delta \cap A$. This is a cocycle, since for every transverse $(d + 1)$ -simplex (Δ', f) the intersection $\Delta' \cap A$ is represented by a compact 1-manifold with boundary and the number of oriented boundary points of such a manifold is 0. If we restrict u to $(\nu^{-1}(x), \nu^{-1}(x) - \{x\})$ for an $x \in A$, we get the orientation class since u sends the generator of the homology to 1. Therefore, u represents the Thom class.

The cellular Gysin map is surely unchanged under subdivisions of the triangulation. Therefore, we can assume that every d -simplex intersects A in at most one point. Furthermore, we can assume via an isotopy of the triangulation that locally at A every d -simplex is only in one fibre. Therefore, we can assume via a suitable subdivision that the p -backface of every $(p + d)$ -dimensional simplex Δ intersecting A maps via the projection map homeomorphically onto $\Delta \cap A$.

A simplex in our triangulation determines a singular simplex with the induced orientation. It is now clear that it is the same if we cap this simplex with u and project it down to A or if we apply the cellular Gysin. This argument clearly works also with local coefficients and proves our claim. This shows in particular that the cellular Gysin does not depend on the chosen triangulation at the level of homology.

3 The Chas-Sullivan Product

In this section, we want to recall the definition of the Chas-Sullivan product. We will follow the approach by David Chataur exhibited in [Cha]. We will fix the notation $L^n M = \text{Map}(S^n, M)$ for the unpointed and $\Omega^n M = \text{Map}^\bullet(S^n, M)$ for the pointed maps.

Let M be an h_* -oriented manifold of dimension d for a homology theory h_* . Consider the diagram

$$\begin{array}{ccc} L^n M \times_M L^n M & \xrightarrow{\iota} & L^n M \times L^n M \\ \text{ev} \downarrow & & \downarrow \text{ev} \times \text{ev} \\ M & \xrightarrow{\Delta} & M \times M \end{array}$$

Here Δ stands for the diagonal, ι is the inclusion and ev the evaluation at the base point pt . Since ev is a submersion, $L^n M \times_M L^n M$ is a sub Hilbert manifold of $L^n M \times L^n M$ and the normal bundle of $L^n M \times_M L^n M$ in $L^n M \times L^n M$ is the pullback of the normal bundle of M in $M \times M$.

We have a map

$$\gamma: L^n M \times_M L^n M = \text{Map}(S^n \vee S^n, M) \rightarrow \text{Map}(S^n, M) = L^n M$$

induced by the collapse map $c: S^n \rightarrow S^n \vee S^n$. The *Chas-Sullivan product* is defined as the composition

$$\begin{array}{ccc}
h_p(L^n M) \otimes h_q(L^n M) & & h_{p+q-d}(L^n M) \\
\downarrow \times & & \uparrow \gamma_* \\
h_{p+q}(L^n M \times L^n M) & \xrightarrow{\iota^!} & h_{p+q-d}(L^n M \times_M L^n M)
\end{array}$$

For notational convenience, we define $\mathbf{h}_*(L^n M) = h_{*+d}(L^n M)$. If $h = H$ is ordinary homology, one usually chooses the notation $\mathbb{H}_*(L^n M) = \mathbf{h}_*(L^n M)$. The above composition may now be written as:

$$\mu: \mathbf{h}_p(L^n M) \otimes \mathbf{h}_q(L^n M) \rightarrow \mathbf{h}_{p+q}(L^n M)$$

Warning 3.1. One has to be careful with signs here since there are different conventions in the literature. The sign convention where the Chas-Sullivan product is graded commutative is used for example in [C-S] and also in [Cha] (where it is ensured by an "artificial" sign), while our sign convention agrees e. g. with that in [C-V] (Theorem 1.2.1 has the wrong sign as stated there).

Remark 3.2. As a aside, we remark that it is also possible to give a completely finite-dimensional bordism-like description of the Chas-Sullivan product in the case of ordinary homology via Kreck's theory of stratifolds. Stratifolds are smooth spaces which are stratified by smooth manifolds satisfying certain conditions on the topology and the relationship of the strata. It is possible to carry out much of the usual differential topology in this setting. Especially we can construct a stratifold bordism homology theory, which coincides with singular homology in the case of spaces having the homotopy type of CW-complexes (for more details and precise definitions see [Kre]). Let now $[S_1, f_1], [S_2, f_2] \in H_*(LM)$ be homology classes. We can interpret a map $f: S \rightarrow LM$ as a map $\text{ev} \circ f: S \rightarrow M$ with a loop in $\Omega(f(p))$ attached to each $p \in S$. Intersect $\text{ev} \circ f_1$ and $\text{ev} \circ f_2$ to get a map $F: S_1 \times_M S_2 \rightarrow M$ and attach to each $p \in S_1 \times_M S_2$ the composition of the loops attached to $\text{pr}_1(p)$ and $\text{pr}_2(p)$. This is a representative of the Chas-Sullivan product of $[S_1, f_1]$ and $[S_2, f_2]$ as is shown in [Mei1, section 3.2.1]. This is close in spirit to the original definition in [C-S].

A closer look at the definition of the Chas-Sullivan product reveals that the only thing we have used of S^n is that it is both a manifold and an H-cogroup (via the map $S^n \rightarrow S^n \vee S^n$). Since every n -dimensional manifold N has the structure of an H-comodule over S^n via the map $c: N \rightarrow N \vee S^n$, collapsing the boundary of a little disc, we get (for M h_* -oriented) the following module structure:

$$\begin{array}{ccc}
h_p(\text{Map}(N, M)) \otimes h_q(L^n M) & & h_{p+q-d}(\text{Map}(N, M)) \\
\downarrow \times & & \uparrow \gamma_* \\
h_{p+q}(\text{Map}(N, M) \times L^n M) & \xrightarrow{\iota^!} & h_{p+q-d}(\text{Map}(N, M) \times_M L^n M)
\end{array}$$

Here $\iota: \text{Map}(N, M) \times_M L^n M \rightarrow \text{Map}(N, M) \times L^n M$ is the inclusion and

$$\gamma: \text{Map}(N, M) \times_M L^n M \rightarrow \text{Map}(N, M)$$

is the map induced by c . This defines a $\mathbf{h}_*(L^n S)$ -module structure on $h_*(\text{Map}(N, M))$, which was first considered in [K-S]. This structure is independent of the chosen disc since

all embeddings of a disc are isotopic. Note that we could substitute $h_p(\text{Map}(N, M))$ by $h'_p(\text{Map}(N, M))$, where h' is a module homology theory over h .

Besides the module structure, there is also the structure of a coalgebra on $\mathbf{h}_*(LM)$ under the condition that $h_*(LM \times_M LM) \cong h_*(LM) \otimes h_*(LM)$ (e.g. for ordinary homology with field coefficients). To define the coproduct, let $i: LM \times_M LM \rightarrow LM$ be the inclusion of all loops $\alpha: [0, 1] \rightarrow M$ with $\alpha(0) = \alpha(\frac{1}{2}) = \alpha(1)$ and $\iota: LM \times_M LM \rightarrow LM \times LM$ the usual inclusion. Then we get a map

$$h_n(LM) \xrightarrow{i^!} h_{n-d}(LM \times_M LM) \xrightarrow{\iota_*} h_{n-d}(LM \times LM) \cong (h_*(LM) \otimes h_*(LM))_{n-d}.$$

This coincides as a special case with the definition of [C-G]. It needs to be said that the coproduct seems all in all to be less interesting than the product structure, at least in the case of singular homology, by recent work of Tamanoi ([Tam]). Note that one cannot mimic the definition above for the higher $L^n M$, because the inclusion $i: L^n M \times_M L^n M \rightarrow L^n M$ has infinite codimension for $n > 1$. Therefore, there are also no interesting comodule structures, since there are not many interesting 1-dimensional manifolds.

4 Spectral Sequences

4.1 Preliminaries

Let $\xi = (E \xrightarrow{\pi} B)$ be a fibre bundle with B a path-connected CW-complex. We define $E^{(p)}$ to be the preimage of the p -skeleton of B under π . Recall that for a homology theory h_* the defining exact couple $C(\xi)$ of the associated Serre spectral sequence is given by

$$\begin{array}{ccc} \bigoplus_{p,q} h_{p+q}(E^{(p)}) & \xrightarrow{i} & \bigoplus_{p,q} h_{p+q}(E^{(p)}) \\ & \swarrow k \quad \searrow j & \\ & \bigoplus_{p,q} h_{p+q}(E^{(p)}, E^{(p-1)}) & \end{array}$$

with the obvious morphisms. The E^1 -term is isomorphic to

$$\bigoplus_{p\text{-cells } \alpha \text{ of } B} [(h_*(D_\alpha^p, S_\alpha^{p-1}) \otimes_{h_*(pt)} h_*(\pi^{-1}(x_\alpha)))]_{p+q},$$

i.e. the p -th part of the cellular complex computing $H_p(B; \mathfrak{h}_q(F))$ where $\mathfrak{h}_*(F)$ denotes the local system given by the homologies of the fibers. It is easy to work out that the isomorphism sends a geometric cycle $[P, a, f]$ to

$$[P \cap \pi^{-1}(x_\alpha), a|_{P \cap \pi^{-1}(x_\alpha)}, f|_{P \cap \pi^{-1}(x_\alpha)}].$$

Here one uses the explicit description of the excision morphism given in 2.2.

To ease the formulation of the results of the next sections, we want to fix some general terminology for spectral sequences. A *morphism of spectral sequences* E_{**}^* and $\tilde{E}_{**}^*$ of level k and bidegree (a, b) consists of homomorphisms $f^n: E_{pq}^n \rightarrow \tilde{E}_{p+a, q+b}^n$ for all $n \geq k$ which commute with the differentials and satisfy $H(f^n) = f^{n+1}$. Now assume that E and $\tilde{E}$

converge to graded abelian groups D_*^∞ and $\tilde{D}_*^\infty$ which are filtered by F_*^* and $\tilde{F}_*^*$ respectively. If we have a morphism f_{**}^* between E and $\tilde{E}$ and in addition homomorphisms $D_r^\infty \rightarrow \tilde{D}_{r+a+b}^\infty$ which map F_r^p to $\tilde{F}_{r+a+b}^{p+a}$ and induce f^∞ on E^∞ , we speak of a *morphism of convergent spectral sequences*. Morphisms of convergent exact couples induce morphisms of the associated convergent spectral sequences.

4.2 Intersecting on Fibre and Base

The goal of this subsection is to define Gysin morphisms of Serre spectral sequences which "compute" the corresponding Gysin morphisms of the homology of the total space.

4.2.1 Intersecting on the Base

Let $\xi = (F \rightarrow E \xrightarrow{\pi} B)$ be a fibre bundle with B a (finite-dimensional) manifold and $A \subset B$ a closed submanifold of codimension d with h_* -oriented normal bundle. Choose a triangulation of B transverse to A and triangulate A as in 2.7.

Theorem 4.1 (Intersection on the Base). *There is a morphism $s_B(A)$ of convergent spectral sequences of level 1 and bidegree $(-d, 0)$ between $\mathcal{E}(\xi)$ and $\mathcal{E}(\xi|_A)$ where the spectral sequences are defined by the triangulations above. The morphism is canonical starting with level 2 and induces the usual Gysin morphism $H_p(B; \mathfrak{h}_q(F)) \rightarrow H_{p-d}(A; \mathfrak{h}_q(F))$ on this level.*

Proof. We want to construct a morphism of the corresponding exact couples $C(\xi)$ and $C(\xi|_A)$. That means, we need to construct morphisms

$$\sigma_a : h_{p+q}(E^{(p)}) \rightarrow h_{p+q-d}(\pi^{-1}(A^{(p-d)}))$$

and

$$\sigma_r : h_{p+q}(E^{(p)}, E^{(p-1)}) \rightarrow h_{p+q-d}(\pi^{-1}(A^{(p-d)}), \pi^{-1}(A^{(p-1-d)}))$$

which commute with the boundary maps. We concentrate on the relative case since this is more difficult.

Let $[P, a, f] \in h_{p+q}(E^{(p)}, E^{(p-1)})$. We want to find a homotopy

$$(P, \partial P) \times I \rightarrow (E^{(p)}, E^{(p-1)})$$

from f to a map g such that πg is transverse to A . To smooth πf , consider open neighbourhoods U_p of the p -skeleton B^p with $U_{p-1} \subset U_p$ such that there are smooth retracts $r_p : U_p \rightarrow B^p$ with $r_p|_{U_{p-1}} = r_{p-1}$. There is a homotopy $H_1 : \partial P \times I \rightarrow U_{p-1}$ from $\pi f|_{\partial P}$ to a smooth map. Extend this homotopy to a homotopy $H_1 : P \times I \rightarrow U_p$ from πf to a map $\tilde{f}$. This map $\tilde{f}$ is smooth on ∂P , so we can find a homotopy $H_2 : P \times I \rightarrow U_p$ to a smooth map such that $H_2|_{\partial P} = \tilde{f} \circ \text{pr}_1$. Now, $r_p \circ H_2(x, 1)$ is homotopic to πf and smooth. Since $E \rightarrow B$ is a fibration we can lift this homotopy. Therefore, we can assume πf to be smooth.

We can homotope $\pi f|_{\partial P}$ in $B^{(p-1)}$ to be transverse to $A \cap B^{(p-1)}$ since $A \cap B^{(p-1)}$ has constant codimension d in $B^{(p-1)}$. We can extend this to a map $\tilde{f}$ on the whole of P . Since $\tilde{f}$ is transverse to A on ∂P , we can homotope it to a $\tilde{g} : P \rightarrow B^{(p)}$ in $B^{(p)}$ which is transverse to $A \cap B^{(p)}$ while leaving ∂P fixed. We can lift this homotopy to E and get a map $g : P \rightarrow E^{(p)}$, for which $\pi g = \tilde{g}$ is transverse to A .

Since $B^p \cap A \subset A^{p-d}$, we can define the maps σ_a and σ_r by transverse intersection of our representative P with A . More precisely, define $Q := (\pi f)^{-1}(A)$ and send $[P, a, f]$ to

$$[Q, a|_Q, f|_Q] \in h_{p+q-d}(E|_{A^{(p-d)}}, E|_{A^{(p-1-d)}})$$

and the same in the absolute case. Since $\partial Q = \partial P \cap Q$, we get a morphism of exact couples and therefore of convergent spectral sequences $E_{**}^1(\xi) \rightarrow E_{(*-d)*}^1(\xi|_A)$.

We now have to check that it induces the usual Gysin morphism on E^2 . To that end, choose the x_α of the last subsection on A for every cell α intersecting A in its interior. Then one sees that our construction coincides with the cellular description of the Gysin morphism. $\square$

Proposition 4.2 (Naturality). *Let $\phi: E' \rightarrow E$ be a map of fibre bundles $\xi' = (F' \rightarrow E' \rightarrow B')$ and $\xi = (F \rightarrow E \rightarrow B)$. Let $A \subset B$ be a submanifold and the map on the bases $f: B' \rightarrow B$ be transverse to A . Then the following diagram commutes beginning with the second level:*

$$\begin{array}{ccc} \mathcal{E}(\xi') & \xrightarrow{\phi_*} & \mathcal{E}(\xi) \\ \downarrow s(f^{-1}(A)) & & \downarrow s(A) \\ \mathcal{E}(\xi'|_{f^{-1}(A)}) & \xrightarrow{\phi_*} & \mathcal{E}(\xi|_A) \end{array}$$

Proof. The normal bundle of $f^{-1}(A)$ in B' is the pullback of the normal bundle of A in B . Since the Thom class is natural, the proposition follows via the Thom isomorphism description of the Gysin morphism. $\square$

We now want to generalize the intersection morphism to an infinite-dimensional context. So let $\xi = (F \rightarrow E \rightarrow B)$ be a fibre bundle with projection map π where B is a Hilbert manifold and $A \subset B$ a closed sub Hilbert manifold of codimension d with h_* -oriented normal bundle. Assume, furthermore, that there is a collection of finite-dimensional manifolds $P_1 \subset P_2 \subset \dots \subset B$ with inclusions $\iota_{i+j}^i: P_i \rightarrow P_{i+j}$ and $\iota_i: P_i \hookrightarrow B$ such that every map $f: X \rightarrow B$ from a compact space can be homotoped into one of the P_i .

Proposition 4.3. *In the situation above, there is a (canonical) morphism $s_B(A)$ of convergent spectral sequences of level 2 and bidegree $(-d, 0)$ between $\mathcal{E}(\xi)$ and $\mathcal{E}(\xi|_A)$. The morphism induces the usual Gysin morphism $H_p(B; \mathfrak{h}_q(F)) \rightarrow H_{p-d}(A; \mathfrak{h}_q(F))$ on E^2 .*

Proof. Let x be in $E_{pq}^n(\xi)$. This element is represented by an element z in $E_{pq}^2 \cong H_p(B; \mathfrak{h}_q(F))$ with $d_2(z) = d_3(z) = \dots = d_{n-1}(z) = 0$. Since

$$H_*(B; \mathfrak{h}_*(F)) \cong \operatorname{colim}_i H_*(P_i; \mathfrak{h}_*(F)),$$

there is an $N \in \mathbb{N}$ such that there is a preimage $y \in E_{pq}^2(\xi|_{P_N})$ of z under the map

$$(\iota_N)_*: \mathcal{E}(\xi|_{P_N}) \rightarrow \mathcal{E}(\xi)$$

with $d_2(y) = d_3(y) = \dots = d_{n-1}(y) = 0$, therefore representing a preimage $[y]$ of x in $E_{pq}^n(\xi|_{P_N})$. By the transversality theorem, we can assume that P_N is transverse to A , hence $P_N \cap A \subset P_N$ is a closed submanifold of P_N . We now define $s_B(A)(x) = (\iota_N)_* s_B(P_N \cap A)([y])$.

We have to check that this is a well-defined map and that it defines a morphism of spectral sequences.

The map is independent of the choices because of the naturality of intersecting on the base: Suppose $(\iota_{n_1})_*(y_1) - (\iota_{n_2})_*(y_2) = d_k(u)$ is a boundary for some $k < n$ where $y_i \in E^2(\xi|_{P_{n_i}})$, $i = 1, 2$, are cycles. We can find a $v \in E^2(\xi|_{P_N})$ ($N \gg 0$) with $\iota_N(v)$ representing u and $\iota_N^{n_1}(y_1) - \iota_N^{n_2}(y_2) = d_k(v)$. Now we use that ι_{n_i} factors over ι_N , that intersecting on the base is natural and that the $\iota_N^{n_1}[y_1] = \iota_N^{n_2}[y_2]$ in $E^n(\xi|_{P_N})$ to deduce that our map is well-defined.

The map is a morphism of spectral sequences since intersecting on the base is a morphism of spectral sequences for each P_i . $\square$

Remark 4.4. If ξ is a smooth fibre bundle of Hilbert manifolds, it is clear by the construction of the intersection on the base that it converges to the usual Gysin morphism on the homology of the total spaces.

4.2.2 Intersecting in the Fibre

Let $\xi = (F \rightarrow E \xrightarrow{\pi} B)$ be a smooth fibre bundle where fibre a Hilbert manifold and base a finite-dimensional manifold. Let $\xi_0 = (F_0 \rightarrow E_0 \rightarrow B)$ be a subbundle of constant codimension d and h_* -oriented normal bundle.

Before we come to our theorem, we want to discuss a variant of the usual description of the Serre spectral sequence. Choose a triangulation on B and consider open neighbourhoods U_p of B^p with $U_{p-1} \subset U_p$ such that there are deformation retracts $r_p: U_p \rightarrow B^p$ with $r_p|_{U_{p-1}} = r_{p-1}$. Then we have an isomorphism from the usual exact couple $C(\xi)$ for the Serre spectral sequence to the exact couple

$$\begin{array}{ccc} \bigoplus_{p,q} h_{p+q}(\pi^{-1}(U_{p-1})) & \xrightarrow{i} & \bigoplus_{p,q} h_{p+q}(\pi^{-1}(U_p)) \\ & \swarrow k \quad \searrow j & \\ & \bigoplus_{p,q} h_{p+q}(\pi^{-1}(U_p), \pi^{-1}(U_{p-1})) & \end{array}$$

Theorem 4.5 (Intersection in the Fibre). *There is a morphism $s_F(E_0)$ of convergent spectral sequences of level 1 and bidegree $(0, -d)$ between $\mathcal{E}(\xi)$ and $\mathcal{E}(\xi_0)$. This induces the usual Gysin morphism $H_p(B; \mathfrak{h}_q(F)) \rightarrow H_p(B; \mathfrak{h}_{q-d}(F_0))$ on E^2 . Furthermore, it converges to the Gysin morphism in the homology of the total spaces.*

Proof. We want to define a morphism of the corresponding exact couples. Let

$$[P, a, f] \in h_{p+q}(\pi^{-1}(U_p), \pi^{-1}(U_{p-1}))$$

be a homology class. As in the proof of 4.1, we can assume f to be smooth.

By proposition 2.2 we can find a homotopy $H: (P, \partial P) \times I \rightarrow (\pi^{-1}(U_p), \pi^{-1}(U_{p-1}))$ from f to a g which is transverse to E_0 . By the same argument, we also can get for a class in $h_{p+q}(\pi^{-1}(U_p))$ representative which is transverse to E_0 . By intersecting with E_0 , we now get a morphism of the exact couples which commutes with the boundary operator as above. More precisely we map $[P, a, f]$ to $[Q, a|_Q, f|_Q] \in h_{p+q-d}(\pi^{-1}(U_p) \cap E_0, \pi^{-1}(U_{p-1}) \cap E_0)$ with $Q = f^{-1}(E_0)$. This induces a convergent morphism $\mathcal{E}(\xi) \rightarrow \mathcal{E}(\xi_0)$ of level 1 and bidegree $(0, -d)$. That this map induces the Gysin morphism on E^2 can be seen by the explicit

isomorphism of the E^1 -term to the cellular complex: there is no difference if we intersect first with F and then with F_0 or if we first intersect with E_0 and then with F_0 (if everything is transverse). $\square$

As in the case of the intersection on the base, we can extend the cases we are interested in to an infinite-dimensional context. So let now B be a Hilbert manifold and the other notation as above and assume that there is a collection of finite-dimensional manifolds $P_1 \subset P_2 \subset \dots \subset B$ such that every map $f: X \rightarrow B$ from a compact space can be homotoped into one of the P_i

Theorem 4.6. *In the situation above, there is a morphism $s_F(E_0)$ of convergent spectral sequences of level 2 and bidegree $(0, -d)$ between $\mathcal{E}(\xi)$ and $\mathcal{E}(\xi_0)$. This induces the usual Gysin morphism $H_p(B; \mathfrak{h}_q(F)) \rightarrow H_p(B; \mathfrak{h}_{q-d}(F_0))$ on E^2 . Furthermore, it converges to the Gysin morphism in the homology of the total spaces.*

Proof. As in the case of the intersection on the base. $\square$

4.3 Multiplicative, Comultiplicative and Module Structures

Define $E[(a, b)]$ to be the shifted spectral sequence with $E^k[(a, b)]_{pq} = E^k_{(p+a)(q+b)}$. For a d -dimensional manifold M , set $\mathbb{H}_*(M) = H_{*+d}(M)$.

Theorem 4.7. *Let M be a d -dimensional h_* -oriented manifold. Then*

$$\mathcal{E}(\Omega^n M \rightarrow L^n M \rightarrow M)[(-d, 0)]$$

can be equipped with the structure of a multiplicative spectral sequence which converges to the Chas-Sullivan product on $\mathfrak{h}_(L^n M)$. Furthermore, the induced product on the E^2 -term $\mathbb{H}_*(M; \mathfrak{h}_*(\Omega^n M))$ is equal to the intersection product with coefficients in the local system of rings $\mathfrak{h}_*(\Omega^n M)$ whose multiplication is given by the Pontryagin product.*

Proof. Let $\xi = (\Omega^n M \rightarrow L^n M \rightarrow M)$ and denote by $\Delta_M: M \rightarrow M \times M$ the diagonal. Then we define the multiplicative structure as the composition

$$\mathcal{E}(\xi) \otimes \mathcal{E}(\xi) \xrightarrow{\times} \mathcal{E}(\xi \times \xi) \xrightarrow{s_B(\Delta_M)} \mathcal{E}(\Delta_M^*(\xi \times \xi)) \xrightarrow{\gamma_*} \mathcal{E}(\xi).$$

Here γ is defined as in 3. Note that the cross product is a map of spectral sequences. All claims are obvious. $\square$

Theorem 4.8. *Let $M \rightarrow N \rightarrow O$ be a fibre bundle of h_* -oriented manifolds of dimensions m, n and o respectively. Then*

$$\mathcal{E}(L^n M \rightarrow L^n N \rightarrow L^n O)[(-o, -m)]$$

can be equipped with the structure of a multiplicative spectral sequence which converges to the Chas-Sullivan product on $\mathfrak{h}_(L^n N)$. Furthermore, the induced product on the E^2 -term $\mathbb{H}_*(L^n O; \mathfrak{h}_{*+m}(L^n M))$ is equal to the Chas-Sullivan product with coefficients in the local system of rings $\mathfrak{h}_{*+m}(L^n M)$.*

Proof. Let $\xi = (L^n M \rightarrow L^n N \rightarrow L^n O)$, $\nu = (L^n M \times_M L^n M \rightarrow L^n N \times_N L^n N \rightarrow L^n O \times_O L^n O)$ and $\iota : L^n O \times_O L^n O \hookrightarrow L^n O \times L^n O$ be the inclusion. We define the multiplicative structure as the composition

$$\mathcal{E}(\xi) \otimes \mathcal{E}(\xi) \xrightarrow{\times} \mathcal{E}(\xi \times \xi) \xrightarrow{s_B(\iota)} \mathcal{E}(\iota^*(\xi \times \xi)) \xrightarrow{s_F(L^n N \times_N L^n N)} \mathcal{E}(\nu) \xrightarrow{\gamma_*} \mathcal{E}(\xi)$$

Here γ is again defined as in section 3. By proposition 2.4, we are in the situation of theorem 4.2.1 and the intersection morphism is defined. $\square$

There is also the notion of a *comultiplicative spectral sequence*, which is simply a comonoid in spectral sequences, i.e. one has a map $E \rightarrow E \otimes E$ which is coassociative (we will not consider counits).

In section 3 we have defined a coproduct on $h_*(LM)$. By the same method, there is also a coproduct on $h_*(\Omega M)$ if we assume h_* to have field coefficients:

$$h_*(\Omega M) \xrightarrow{i^!} h_{*-d}(\Omega M \times \Omega M) \cong [h_*(\Omega M) \otimes_{h_*} h_*(\Omega M)]_{*-d}$$

Here $i : \Omega M \times \Omega M \rightarrow \Omega M$ is the inclusion of loops α with $\alpha(\frac{1}{2}) = \alpha(0)$. For V a coalgebra over a field k , we have furthermore a coproduct on $H_*(M; V)$:

$$H_*(M; V) \rightarrow H_*(M; V \otimes V) \xrightarrow{\Delta_*} H_*(M \times M; V \otimes V) \cong H_*(M; V) \otimes_k H_*(M; V)$$

Theorem 4.9. *Let M be a d -dimensional h_* -oriented manifold with h_* a field. Then*

$$\mathcal{E}(\Omega M \rightarrow LM \rightarrow M)[(0, -d)]$$

can be equipped with the structure of a comultiplicative spectral sequence which converges to the coproduct on $h_(LM)$. Furthermore, the induced coproduct on the E^2 -term $H_*(M; \mathfrak{h}_{*+d}(\Omega M))$ is equal to the coproduct on M with coefficients in the coalgebra $\mathfrak{h}_{*+d}(\Omega M)$.*

Proof. Let ξ be as above and $\xi \times_M \xi = (\Omega M \times \Omega M \rightarrow LM \times_M LM \rightarrow M)$. Consider the diagram

$$\mathcal{E}(\xi) \xrightarrow{s_F(LM \times_M LM)} \mathcal{E}(\xi \times_M \xi) \xrightarrow{\Delta_*} \mathcal{E}(\xi \times \xi) \xrightarrow{\cong} \mathcal{E}(\xi) \otimes \mathcal{E}(\xi)$$

All claims are obvious. $\square$

Theorem 4.10. *Let $M \rightarrow N \rightarrow O$ be a fibre bundle of h_* -oriented manifolds of dimensions m , n and o , respectively with h_* a field. Then*

$$\mathcal{E}(LM \rightarrow LN \rightarrow LO)[(-o, -m)]$$

can be equipped with the structure of a comultiplicative spectral sequence which converges to the coproduct on $h_(LN)$. Furthermore, the induced coproduct on the E^2 -term $\mathbb{H}_*(LO; \mathfrak{h}_{q+m}(LM))$ is equal to the coproduct with coefficients in the coalgebra $\mathfrak{h}_{*+m}(LO)$.*

Proof. Analogous to the previous theorem and the one for the corresponding multiplicative spectral sequence. $\square$

For E a multiplicative spectral sequence, there is also the notion of a module spectral sequence, i.e. a spectral sequence E' together with a morphism $E \otimes E' \rightarrow E'$ and the usual coherence diagrams. Recall that, if M is a d -manifold and N a module over a ring R , we have an $\mathbb{H}_*(M; R)$ -module structure on $\mathbb{H}_*(M; N)$ defined analogous to the intersection product (here we use the cross product $H_*(M; R) \otimes H_*(M; N) \rightarrow H_*(M \times M; N)$).

Theorem 4.11. *Let Z be a closed n -manifold and M be a d -dimensional h_* -oriented manifold. Then*

$$\mathcal{E}(\mathrm{Map}^\bullet(Z, M) \rightarrow \mathrm{Map}(Z, M) \rightarrow M)[(-d, 0)]$$

can be equipped with the structure of a module spectral sequence over $\mathcal{E}(\Omega^n M \rightarrow L^n M \rightarrow M)[(-d, 0)]$ which converges to the module structure on $h_(\mathrm{Map}(Z, M))$. Furthermore, the induced module structure on the E^2 -term $\mathbb{H}(M; h_*(\mathrm{Map}^\bullet(Z, M)))$ coincides with the module structure described above.*

Proof. As in the multiplicative case. □

Theorem 4.12. *Let Z be a closed n -manifold. Furthermore, let $M \rightarrow N \rightarrow O$ be a fibre bundle of h_* -oriented manifolds of dimensions m , n and o respectively. Then*

$$\mathcal{E}(\mathrm{Map}(Z, M) \rightarrow \mathrm{Map}(Z, N) \rightarrow \mathrm{Map}(Z, O))[-o, -m]$$

can be equipped with the structure of a module spectral sequence over $\mathcal{E}(L^n M \rightarrow L^n N \rightarrow L^n O)[-o, -m]$ which converges to the module structure on $h_(\mathrm{Map}(Z, N))$. Furthermore the induced module structure on the E^2 -term*

$$H_{*+o}(\mathrm{Map}(Z, O); h_{*+m}(\mathrm{Map}(Z, M)))$$

coincides with the module structure described in 3.

Proof. As in the multiplicative case. □

5 Examples

In this subsection, we will do two different things, in object and method. First, we want to widen our knowledge about ordinary homology of free loop spaces to the case of certain sphere bundles. Secondly, we want to compute some extraordinary homologies of free loop spaces, namely Landweber exact theories (e.g. complex cobordism and complex K-Theory) and oriented bordism. We will study the Atiyah-Hirzebruch spectral sequence associated to spheres and (complex) projective space and show that it degenerates on E^2 . To achieve this, we need to construct first explicit manifold generators for the ordinary homology of the free loop spaces of the spheres and projective spaces, which may be interesting in its own right.

5.1 The Case of Sphere Bundles

We want to study the homology of the free loop space of sphere bundles. While the integral homology of free loop spaces is usually hard to compute, there are more efficient tools for the rational homology, namely rational homotopy theory. Recall that rational homotopy theory associates functorially to every simply-connected space X a (graded) commutative differential graded algebra (dga) over $\mathbb{Q}$ of the form ΛV . Here V is a graded rational vector space and ΛV is the free (graded) commutative dga over V , i.e. a tensor product of polynomial rings for the basis elements of V of even degree and exterior algebras for the odd parts. This is called the *minimal model* $\mathcal{M}(X)$ of X . The cohomology of $\mathcal{M}(X) = \Lambda V$ is isomorphic to the rational cohomology of X (see [FHT]).

We will need the following two facts of rational homotopy theory:

1. The vector space V is naturally isomorphic to the dual of $\pi_*(X; \mathbb{Q}) := \pi_*(X) \otimes \mathbb{Q}$ (see [FHT], Thm 15.11).
2. The minimal model of LX depends only on the minimal model of X . This can be seen by the explicit formulas of [V-S].

While the minimal model of LX only gives information about the rational cohomology, we want to use rational homotopy theory in combination with the Serre spectral sequence to do integral computations for the free loop space LE of a fibre bundle $S^k \rightarrow E \rightarrow S^n$. We use the computation of $\mathbb{H}(S^n)$ for $n > 1$ by Cohen, Jones and Yan ([CJY]):

$$\begin{aligned}\mathbb{H}_*(LS^n) &= \Lambda(a) \otimes \mathbb{Z}[u] \text{ for } n \text{ odd,} \\ \mathbb{H}_*(LS^n) &= \Lambda(b) \otimes \mathbb{Z}[a, v]/(a^2, ab, 2av) \text{ for } n \text{ even}\end{aligned}$$

with generators $a \in \mathbb{H}_{-n}(LS^n)$, $b \in \mathbb{H}_{-1}(LS^n)$, $u \in \mathbb{H}_{n-1}(LS^n)$ and $v \in \mathbb{H}_{2n-2}(LS^n)$. For this computation they use the multiplicative spectral sequence exhibited in 4.3 in the special case of singular homology.

First assume $k > 1, n > 1$ odd. The odd dimensional spheres have only one nontrivial rational homotopy group, namely $\pi_k(S^k; \mathbb{Q}) = \mathbb{Q}$. Hence $\pi_i(E; \mathbb{Q}) = \pi_i(S^k; \mathbb{Q}) \oplus \pi_i(S^n; \mathbb{Q})$ in every degree by the long exact sequence of homotopy groups. So we have $\mathcal{M}(E) = \Lambda(x_k) \otimes \Lambda(x_n)$ with $|x_k| = k$ and $|x_n| = n$. For dimension reasons, there are no differentials. Thus we have $\mathcal{M}(E) \cong \mathcal{M}(S^k \times S^n)$ as differential graded algebras. We conclude

$$H_*(LE; \mathbb{Q}) \cong H_*(L(S^k \times S^n); \mathbb{Q}).$$

Consider the E^2 -term of the Serre spectral sequence associated to $LS^k \rightarrow LE \rightarrow LS^n$. Every occuring group is torsionfree. Therefore, our rational computation shows that the spectral sequence degenerates at E^2 and we have

$$\mathbb{H}_*(LE) \cong \mathbb{H}_*(LS^k) \otimes \mathbb{H}_*(LS^n) \cong \Lambda(a_k, a_n) \otimes \mathbb{Z}[u_k, u_n]$$

with $|a_k| = -k$, $|a_n| = -n$, $|u_k| = k - 1$ and $|u_n| = n - 1$ (for the grading conventions, see section 3). Note that all extension are trivial in the sense that

$$0 \rightarrow F^{n-1} \rightarrow F^n \rightarrow F^n/F^{n-1} \rightarrow 0$$

splits since all occuring groups in E^∞ are torsionfree. To show that the isomorphism above holds also multiplicatively, we apply the following proposition to the Serre spectral sequence associated to $LM \rightarrow LN \rightarrow LO$:

Proposition 5.1. *Let E be a multiplicative convergent first quadrant spectral sequence of modules over a noetherian ring R that converges multiplicatively to a graded group G_* and has the grading conventions of the homological Serre spectral sequence. Assume that $E_{**}^\infty \cong E_{*0}^\infty \otimes E_{0*}^\infty$ and that this is finitely generated over R in every bidegree. Furthermore, require that all extension are trivial and $E_{*0}^\infty = R[\bar{x}_0, \bar{x}_1, \dots, \bar{x}_n] \otimes \Lambda_R(\bar{x}_{n+1}, \bar{x}_{n+2}, \dots)$. Then $G_k \cong \bigoplus_{p+q=k} E_{pq}^\infty$ holds multiplicatively.*

Proof. Denote the filtration of F_q by F_q^* . As $E_{p0}^\infty = F_p^p/F_p^{p-1}$ and $F_p^p = G_p$, we have a surjective map $G_p \rightarrow E_{p0}^\infty = E_{p0}^2$. Lift the $\bar{x}_j$ to x_j in G_p . Since the multiplication on

$E^\infty = F^p/F^{p-1}$ is induced by that on G_* , we have that $x_i x_j$ is a lift for $\overline{x_i x_j}$. The groups E_{0*}^∞ act on G_* and E^∞ in a compatible way via multiplication. Consider the map

$$L: E_{**}^\infty \rightarrow G_*, \quad y \cdot \Pi x_i^{k_i} \mapsto y \cdot \Pi \overline{x_i}^{k_i},$$

where $y \in E_{0*}^\infty$. This map is clearly a map of algebras. It is also clear that it is surjective onto F^0 . Assume inductively that it is surjective onto F^p . The products $\Pi \overline{x_i}^{k_i}$ with $\sum k_i |x_i| = p+1$ form a E_{0*}^∞ -basis for F^{p+1}/F^p and are images of L . Therefore, we see that L is surjective onto F^{p+1} and conclude by induction that it is surjective onto the whole of G_* . Since $E_{**}^\infty \cong G_*$ additively and both are finitely generated (hence noetherian) R -modules in every degree, L is an isomorphism (of algebras). Indeed, identify E_{**}^∞ and G_* and consider the ascending chain $\ker(L^n)$, $n \in \mathbb{N}$. For some N , we have $\ker(L^N) = \ker(L^{N+1})$. Since every $y \in G_*$ is of the form $L^N(y')$, we get $Ly = 0$ iff $y = 0$. $\square$

Now we consider the case $k > 1$ odd and $n > 2$ even. Assume furthermore that $k \neq n \pm 1$ and that $n - 1$ is no multiple of $k - 1$. Even dimensional spheres S^n have two non-zero rational homotopy groups, namely $\pi_n(S^n; \mathbb{Q}) \cong \mathbb{Q}$ and $\pi_{2n-1}(S^n, \mathbb{Q}) \cong \mathbb{Q}$. By the long exact homotopy sequence, we have

$$\pi_i(E; \mathbb{Q}) = \pi_i(S^k; \mathbb{Q}) \oplus \pi_i(S^n; \mathbb{Q})$$

in every degree. So we get

$$\mathcal{M}(E) \cong \Lambda(x_k) \otimes \mathbb{Z}[x_n] \otimes \Lambda(y_{2n-1})$$

with $|x_k| = k$, $|x_n| = n$ and $|y_{2n-1}| = 2n - 1$. Since the Serre spectral sequence associated to $S^k \rightarrow E \rightarrow S^n$ degenerates at E^2 , we have $d(x_k) = d(x_n) = 0$ and $d(y_{2n-1})$ must be a non-zero multiple of x_n^2 . Therefore, $\mathcal{M}(E)$ is isomorphic to the minimal model $\mathcal{M}(S^k \times S^n)$ and hence we have

$$H_*(LE; \mathbb{Q}) \cong H_*(LS^k \times LS^n; \mathbb{Q}).$$

Consider the Serre spectral sequence associated to $LS^k \rightarrow LE \rightarrow LS^n$. A differential $d_i(x)$ can only be non-zero if $d_i(x)$ is torsion. The only torsion elements of $\mathbb{H}_*(LS^n)$ are the av^j for $j \geq 1$ (see 5.2.1 for notation). Hence, we have

$$d_i(1 \otimes a_{LS^k}) = d_i(1 \otimes u_{LS^k}) = d_i(a_{LS^n} \otimes 1) = d_i(b_{LS^n} \otimes 1) = d_i(v_{LS^n} \otimes 1) = 0$$

for all $i \geq 2$ as one sees by an analysis of possible differentials. The E^2 -term of the Serre spectral sequence is isomorphic to $\mathbb{H}_*(LS^n) \otimes \mathbb{H}_*(LS^k)$. By multiplicativity, the spectral sequence degenerates at E^2 . Because filtration issues may come up, we cannot deduce in this case the concrete structure of the homology.

5.2 Manifold Generators

5.2.1 The Spheres

In this section, we will present concrete generators for the homology of LS^n for $n > 1$. To achieve this, we consider first the simpler case of ΩS^n . It is well known that $H_*(\Omega S^n) \cong \mathbb{Z}[x]$ with $x \in H_{n-1}(\Omega S^n)$, where the product is induced by composing loops. By adjunction from the identity, we get a map $f: S^{n-1} \rightarrow \Omega \Sigma S^{n-1} \cong \Omega S^n$. This represents a class in $H_{n-1}(\Omega S^n)$, which is easy to be seen a additive generator.

To visualize x , think of the base point as the north pole. The points $p \in S^{n-1}$ of the equator parametrize the minimal geodesics γ_p between north and south pole. Now choose a distinguished minimal geodesic δ from the south to the north pole (the "way backwards"). Then $p \mapsto \delta * \gamma_p$ defines $f: S^{n-1} \rightarrow \Omega S^n$ where $*$ denotes the concatenation of paths (note that the suspensions above are reduced).

Now consider the free loop space. We use the same notation for the generators of homology as in the last section. Since a is in $H_0(LM)$, it can be represented by an arbitrary loop, e.g. a constant loop. By studying the Serre spectral sequence for $\Omega M \rightarrow LM \rightarrow M$, it can be shown that $j_*: H_{n-1}(\Omega M) \rightarrow \mathbb{H}_{-1}(LM)$ is an isomorphism (see [CJY]). Therefore, $j_*(x)$ is a generator of $\mathbb{H}_{-1}(LM)$ and hence up to sign equal to b for n even and to au for n odd.

For identifying the other generators, we begin with the easier case of n odd. Consider the Gysin morphism

$$j^!: H_*(LS^n) \rightarrow H_{*-d}(\Omega S^n).$$

We want to find a preimage of x under $j^!$. This is a fortiori a generator of $\mathbb{H}_{n-1}(LS^n)$ and therefore up to sign equal to u . Let S^n be equipped with the standard metric of the sphere of circumference 1 and STS^n be the unit sphere bundle in the tangent bundle TS^n . Let V be a vector field of unit length. We define a map $F: STS^n \rightarrow LS^n$ by

$$(p, v) \mapsto \left(t \mapsto \begin{cases} \exp_p(tv) & \text{for } t \leq \frac{1}{2} \\ \exp_{-p}((t - \frac{1}{2})V(-p)) & \text{for } t \geq \frac{1}{2} \end{cases} \right)$$

Here p denotes a point in S^n and v is a unit tangent vector to p . By the description of x above it is clear that $j^!(F_*[STS^n]) = x$.

This construction cannot work for n even, since in this case we have $\mathbb{H}_{n-1}(LS^n) = 0$ for $n > 2$ and the generator $bv \in \mathbb{H}_{n-1}(LS^2)$ maps to zero under $j^!$, because a representative (S, f) can be chosen with $\text{im}(\text{ev} \circ f) = pt$.³

To construct an explicit representative of v , we need an alternative representative of x^2 . By our description above we get as a representative:

$$(v_1, v_2) \mapsto \left(s \mapsto \begin{cases} \exp_p(2sv_1) & \text{for } s \leq \frac{1}{4} \\ \exp_{-p}(2(s - \frac{1}{4})w) & \text{for } \frac{1}{4} \leq s \leq \frac{1}{2} \\ \exp_p(2(s - \frac{1}{2})v_2) & \text{for } \frac{1}{2} \leq s \leq \frac{3}{4} \\ \exp_{-p}(2(s - \frac{3}{4})(-w)) & \text{for } \frac{3}{4} \leq s \leq 1 \end{cases} \right)$$

Here v_1 and v_2 are unit tangent vectors at the base point p and w is a unit tangent vector at $-p$. This map is now easy to be seen to be homotopic to

$$(v_1, v_2) \mapsto \left(s \mapsto \begin{cases} \exp_p(sv_1) & \text{for } s \leq \frac{1}{2} \\ \exp_{-p}((s - \frac{1}{2})(-v_2)) & \text{for } \frac{1}{2} \leq s \leq 1 \end{cases} \right) \quad (1)$$

where $-v_2$ denotes the parallel transport of v_2 along any geodesic from p to $-p$. By this description, it is now easy to construct a preimage of x^2 under $j^!$: consider the pullback

³This gives an eccentric proof for the theorem of the hairy ball, because the only thing we used for n odd was the existence of a non-vanishing vector field.

E of the product bundle $STS^n \times STS^n$ via the diagonal $S^n \rightarrow S^n \times S^n$. Then construct $F: E \rightarrow LS^n$ by defining $F(p, v_1, v_2)$ by the formula above (1). This is obviously a preimage of x^2 . Since x^2 is an additive generator of $H_{2n-2}(\Omega S^n)$ and $\mathbb{Z}\{v\}$ is the non-torsion part of $\mathbb{H}_{2n-2}$, the class $[E, F]$ equals v up to sign.

By the Chas-Sullivan product we get now explicit generators for every class in $H_*(LM)$. In particular every class in $H_*(LM)$ is represented by a manifold.

Remark 5.2. We could also have used the explicit generators to deduce the multiplicative structure of $H_*(LM)$ from the additive structure, without using the compatibility of the Chas-Sullivan product with the Serre spectral sequence.

5.2.2 Projective Spaces

In [CJY] the authors show that

$$\mathbb{H}_*(L\mathbb{CP}^n) \cong \Lambda(w) \otimes \mathbb{Z}[c, u]/(c^{n+1}, wc^n, (n+1)c^nu),$$

where $|w| = -1$, $|c| = -2$ and $|u| = 2n$. By the same methods one can show

$$\mathbb{H}_*(L\mathbb{HP}^n) \cong \Lambda(w) \otimes \mathbb{Z}[c, u]/(c^{n+1}, wc^n, (n+1)c^nu)$$

with $|w| = -1$, $|c| = -4$ and $|u| = 4n+2$. We will find explicit generators for these homology classes. In the following, $\mathbb{K}$ will stand for one of $\mathbb{C}$ or $\mathbb{H}$ and d will be the $\mathbb{R}$ -dimension of $\mathbb{K}$.

There is a homotopy fibration $S^{d(n+1)-1} \rightarrow \mathbb{KP}^n \rightarrow \mathbb{KP}^\infty$ (see [CJY]). If we loop this, we get a homotopy section $\Omega(\mathbb{KP}^\infty) \rightarrow \Omega(\mathbb{KP}^n)$, since the map $\Omega\mathbb{KP}^\infty \simeq S^{d-1} \rightarrow S^{d(n+1)-1}$ is nullhomotopic. Therefore we get (additively):

$$H_*(\Omega\mathbb{KP}^n) \cong H_*(\Omega S^{d(n+1)-1}) \otimes H_*(S^{d-1}) \cong \mathbb{Z}[y] \otimes \Lambda(z),$$

where $|y| = d(n+1)-2$ and $|z| = d-1$. Here we use $\Omega\mathbb{KP}^\infty \simeq S^{d-1}$. The above isomorphism holds also multiplicatively, because the Serre spectral sequence is here multiplicative, since the Pontrjagin product is induced by a map and the Serre spectral sequence is natural (note that there are no filtration issues for dimension reasons).

In this whole section, let the projective spaces be equipped with the metric coming from the standard metric on the unit sphere. The generator c is represented by $c: \mathbb{KP}^{n-1} \hookrightarrow \mathbb{KP}^n \hookrightarrow L\mathbb{KP}^n$. This holds because $\text{ev} \circ c: \mathbb{KP}^{n-1} \rightarrow \mathbb{KP}^n$ represents a generator of $\mathbb{H}_{-d}(\mathbb{KP}^n)$.

Let $d > 1$ and $STS^{d(n+1)-1}$ be the sphere bundle of the tangent bundle of $S^{d(n+1)-1}$. Let E be the quotient of the sphere bundle via the (isometric) action of S^{d-1} on $S^{d(n+1)-1}$. Clearly the composite $STS^{d(n+1)-1} \rightarrow LS^{d(n+1)-1} \rightarrow L\mathbb{KP}^n$ of the generator u of $H_*(LS^{d(n+1)-1})$ and the looped Hopf map factors over E . This map $E \rightarrow L\mathbb{KP}^n$ is our generator u . To see this, we simply intersect u with $\Omega\mathbb{KP}^n$ and observe that this equals y in the notation above.

For the last generator look at $T\mathbb{KP}^n|_{\mathbb{KP}^{n-1}}$. This has certainly a nonvanishing section s by obstruction theory, because the (dn) -th cohomology of $\mathbb{KP}^{n-1}$ vanishes. We can assume that $s(x)$ has unit length for every $x \in \mathbb{KP}^{n-1}$. Let now L be the (trivial) S^{d-1} -bundle in $T\mathbb{KP}^n|_{\mathbb{KP}^{n-1}}$ generated by s . Then we define our map $w': L \rightarrow L\mathbb{KP}^n$ via

$$(p, l) \mapsto \begin{cases} t \mapsto \exp_p(\pi t \cdot l) & \text{for } t \leq \frac{1}{2} \\ t \mapsto \exp_p(\pi(1-t) \cdot s(p)) & \text{for } t \geq \frac{1}{2} \end{cases}$$

If we multiply the represented homology class $[w']$ with c^{n-1} , i.e. intersect with a $\mathbb{K}\mathbb{P}^1$ connecting a $p \in \mathbb{K}\mathbb{P}^{n-1}$ with $\exp_p(\frac{\pi}{2} \cdot s(p))$, we get obviously the image of the generator $[f] \in H_{d-1}(\Omega S^d) \cong H_{d-1}(\Omega \mathbb{K}\mathbb{P}^1)$ under the map $\Omega \mathbb{K}\mathbb{P}^1 \hookrightarrow \Omega \mathbb{K}\mathbb{P}^n \hookrightarrow L\mathbb{K}\mathbb{P}^n$. The first is an isomorphism on H_{d-1} because of the description of the homology of $\Omega \mathbb{K}\mathbb{P}^n$ above. The latter is also an isomorphism on H_1 by inspection of the Serre spectral sequence associated to $\Omega \mathbb{K}\mathbb{P}^n \rightarrow L\mathbb{K}\mathbb{P}^n \rightarrow \mathbb{K}\mathbb{P}^n$ (see [CJY]). So w' is an additive (non-torsion) generator since $[w']c^{n-1}$ is. Since $\mathbb{H}_{-1}(L\mathbb{K}\mathbb{P}^n) \cong \mathbb{Z}$, this settles w' (modulo sign) as the generator w described in [CJY].

Remark 5.3. A similar description in the case $\mathbb{K} = \mathbb{R}$ can be found in [Mei1, section 3.6.2].

5.3 Calculations in Generalized Homology Theories

5.3.1 Complex Cobordism and Landweber Exact Theories

We define natural transformations $\mu : MSO_n(X) \rightarrow H_n(X)$ and $\nu : MU_n(X) \rightarrow H_n(X)$ by sending a representing cycle $[M, f]$ to $f_*([M])$.

We have the following well-known proposition:

Proposition 5.4. *If X is (homotopy equivalent to) a CW-complex, then the MU spectral sequence degenerates at E^2 if and only if $\nu : MU_n(X) \rightarrow H_n(X)$ is an epimorphism for all $n \geq 0$.*

To see the degeneration of the Atiyah-Hirzebruch spectral sequence for MU , we have to find stably almost complex structures on our generators of 5.2.1 and 5.2.2. Since S^n has trivial normal bundle in $\mathbb{R}^{n+1}$, we have that $TS^n \oplus \varepsilon$ is trivial, where ε is the trivial line bundle. Since trivial bundles are stably complex, the sphere is stably almost complex. Recall, we denoted the sphere subbundle of the tangent bundle by STS^n . We have that $STS^n \subset S(TS^n \oplus \varepsilon) \cong S^n \times S^n$ has trivial normal bundle. Since the tangent bundle of $S^n \times S^n$ is stably trivial, the tangent bundle of STS^n is stably trivial, too, and therefore stably complex. This finishes the case for the sphere.

The manifolds $\mathbb{C}\mathbb{P}^n$ and $\mathbb{C}\mathbb{P}^n \times S^1$ are clearly almost complex. It remains to show that STS^{2n+1}/S^1 is stably almost complex, where S^1 acts via complex multiplication and its derivative. As in the paragraph above, it suffices to consider $(S^{2n+1} \times S^{2n+1})/S^1$, where S^1 acts via the diagonal action. Embed $S^{2n+1} \times S^{2n+1}$ into $\mathbb{C}^{n+1} \times \mathbb{C}^{n+1}$. This gives an embedding $(S^{2n+1} \times S^{2n+1})/S^1 \hookrightarrow (\mathbb{C}^{n+1} \times \mathbb{C}^{n+1})/\mathbb{C}^* \cong \mathbb{C}\mathbb{P}^{2n+1}$ of codimension 1. Since the latter is complex and the normal bundle is trivial (note that $(S^{2n+1} \times S^{2n+1})/S^1$ is simply connected), degeneration is proven.

By Localizing at p , we can deduce that the Atiyah-Hirzebruch spectral sequence for free loop spaces of spheres and complex projective spaces degenerates at E^2 also for BP . It is not difficult to show that one gets thereby degeneration also for all Landweber exact theories, i.e. homology theories h_* of the form $h_*(X) = MU_*(X) \otimes_{MU_*} h_*$ or $h_*(X) = BP_*(X) \otimes_{BP_*} h_*$. This includes, among others, complex K -homology, elliptic homology, the Johnson-Wilson theories $E(n)$ and the Morava E -theories E_n . Note that the isomorphism $h_*(LM) = MU_*(LM) \otimes_{MU_*} h_*$ or $h_*(LM) = BP_*(LM) \otimes_{BP_*} h_*$ holds multiplicatively with respect to the Chas-Sullivan product for a MU_* - respectively BP_* -oriented manifold M .

For the odd dimensional spheres, we have in addition that all filtration extension are trivial, i.e.

$$0 \rightarrow F^{n-1} \rightarrow F^n \rightarrow F^n/F^{n-1} \rightarrow 0$$

splits since F^n/F^{n-1} is a free h_* -module. Therefore, we have additively

$$h_*(LS^{2k+1}) \cong H_*(LS^{2k+1}) \otimes h_*$$

for every Landweber exact theory h_* .

We have $E^2(LM, MU)_{pq} = H_p(LM) \otimes MU_*(pt)$. By 5.1, the above isomorphisms

$$\begin{aligned} MU_*(LS^{2k+1}) &\cong H_*(LS^{2k+1}) \otimes MU_* \\ BP_*(LS^{2k+1}) &\cong H_*(LS^{2k+1}) \otimes BP_* \end{aligned}$$

hold now also multiplicatively. Thereby, we can conclude an analogous isomorphism for every Landweber exact homology theory.

In the light of the physical interest in equivariant index theory on the free loop space, it would be exciting to extend these calculations to the equivariant K -theory of free loop spaces.

5.3.2 Oriented Bordism

As in 5.3.1, we have

Proposition 5.5 ([C-F], 15.1). *If X is (homotopy equivalent to) a CW-complex then the Atiyah-Hirzebruch spectral sequence for oriented bordism degenerates at E^2 if and only if $\mu: MSO_n(X) \rightarrow H_n(X)$ is surjective for all $n \geq 0$.*

As we have described concrete manifold generators in 5.2.1 and 5.2.2, we get degeneration for free loop spaces of spheres and (complex and quaternionic) projective spaces. But we can prove even more in some cases:

Theorem 5.6 ([C-F], 15.2). *If X is (homotopy equivalent to) a CW-complex for which each $H_n(X)$ is finitely generated and has no odd torsion, then*

$$MSO_n(X) \cong \bigoplus_{p+q=n} H_p(X; MSO_q)$$

We can apply this theorem to these free loop spaces of spheres and complex or quaternionic projective spaces which have no odd torsion in homology. Therefore, we have additive isomorphisms

$$\begin{aligned} MSO_*(LS^{2k}) &\cong H_*(LS^{2k}; MSO_*(pt)) \\ MSO_*(LS^{2k+1}) &\cong H_*(LS^{2k+1}) \otimes MSO_*(pt) \\ MSO_*(\mathbb{CP}^{2^k-1}) &\cong H_*(L\mathbb{CP}^{2^k-1}; MSO_*(pt)) \\ MSO_*(\mathbb{HP}^{2^k-1}) &\cong H_*(L\mathbb{HP}^{2^k-1}; MSO_*(pt)) \end{aligned}$$

Sadly enough, MSO_* is not torsionfree, but has also 2-torsion (a complete determination can be found in [Wal]). Therefore, we cannot decide by this method whether these isomorphisms hold also for the multiplicative structure except in the case of odd-dimensional spheres, where the above isomorphism does hold multiplicatively.

References

- [C-S] M. Chas, D. Sullivan, *String Topology*, arXiv:math/0105178v2 (1999)
- [Cha] D. Chataur, *A Bordism Approach to String Topology*, math.AT/0306080 (2003)
- [C-G] R.L. Cohen, V. Godin, *A polarized view of string topology* in *Topology, Geometry, and Quantum Field theory*, Lond. Math. Soc. lecture notes vol. 308 (2004), 127-154.
- [C-J] R.L. Cohen, J.D.S Jones, *A homotopy theoretic realization of string topology* in *String Topology and Cyclic Homology*, pp. 1-96, Birkhäuser (2006). Online available at arXiv:math/0107187v2
- [C-V] R.L. Cohen, A. Voronov, *Notes on String Topology*, arXiv:math/0503625v1
- [CJY] R.L. Cohen, J.D.S. Jones und J. Yan, *The loop homology algebra of spheres and projective spaces*, Progr. Math., Vol. 215, Birkhäuser, Basel, 2003, 77-92
- [C-F] P. E. Conner, *Differentiable Periodic Maps*, Springer (1970)
- [FHT] Y. Félix, S. Halperin, J-C Thomas, *Rational Homotopy Theory*, Springer (2001)
- [Jak] M. Jakob, *An Alternative Approach to Homology*, Contemporary Mathematics 265 (2000), 87-97
- [K-S] S. Kallel, P. Salvatore, *Rational maps and string topology*, Geom. Topol. 10 (2006) 1579-1606
- [Kre] M. Kreck, *Differential Algebraic Topology*, http://www.mathi.uni-heidelberg.de/~kreck/stratifolds/DA_ (2005)
- [LBo] J-F Le Borgne, *The loop-product spectral sequence*, Expositiones Mathematicae, Vol 26, Issue 1 (2008), 25-40
- [LBo2] J-F Le Borgne, *The loop-coproduct spectral sequence*, arXiv:math/0512276v1 (2005)
- [Mei1] L. Meier, *A Geometric View on String Topology*, Diplomarbeit (2009). Available at the author's homepage <http://www.math.uni-bonn.de/people/lmeier>
- [Mei2] L. Meier, *A Hilbert Manifold Model for Mapping Spaces*. Available at the author's homepage <http://www.math.uni-bonn.de/people/lmeier>
- [Mil] J. Milnor, *Morse Theory*, Princeton University Press (1963)
- [Qui] F. Quinn, *Transversal approximation on Banach manifolds*, Proc. Symp. Pure Math. 15 (1970), 213-222.
- [Tam] H. Tamanoi, *Loop Coproducts in String Topology and Triviality of Higher Genus TQFT Operations*, arXiv:0706.1276v3
- [V-S] M. Vigue-Poirrier, D. Sullivan, *The Homology Theory of the Closed Geodesic Problem*, J. Differential Geometry 11 (1976), 633-644
- [Wal] C.T.C. Wall, *Determination of the Cobordism Ring*, Annals of Mathematics, Vol. 72, No. 2 (Sept. 1960), 292-311.